

ITFA 93-05
 ITEP-M-1/93
 hep-th/9303150

Charge screening in the Higgs phase of Chern-Simons electrodynamics

F.Alexander Bais*, Alexei Morozov^{†,‡}, and Mark de Wild Propitius[§]

Institute for Theoretical Physics
 Valckenierstraat 65, 1018XE Amsterdam, the Netherlands

March 1993

Abstract

Though screened at large distances, a point-like electric charge can still participate in a long-range electromagnetic interaction in the Higgs phase, namely that with the Aharonov-Bohm field produced by a localized magnetic flux. We show that this follows from the fact that the screening charge, induced in the presence of a Higgs condensate, does not interact with the Aharonov-Bohm field. The same phenomenon occurs, if a Chern-Simons term is incorporated in the action. This observation provides a physical basis for the recently proposed classification of the superselection sectors of this model in terms of a quasi-Hopf algebra.

PACS numbers: 0365, 0530L, 1130, 1220

*bais@phys.uva.nl

[†]Permanent address: ITEP 117259, Moscow, Russia

[‡]morozov@vxdesy.desy.de; morosov@phys.uva.nl

[§]mdwp@phys.uva.nl

The question whether charges in the Higgs phase can be measured through Aharonov-Bohm (AB) scattering with magnetic fluxes [1], has received quite some attention in recent literature. These discussions mostly focussed on gedanken experiments involving closed shells or annuli of unbroken vacuum surrounding the charge (see for instance Ref. [2]), and lead to the conclusion that Coulomb fields need not be screened inside the volume V surrounded by these shells (i.e. $Q_V = \int_V \partial_i F^{i0} d^D x \neq 0$). The point we like to make in this letter, is that even *inside* the broken phase these charges can be measured by means of AB scattering with magnetic fluxes (and vice versa) due to a rather subtle effect, which depends on the specific details of the Higgs mechanism. So, though the Coulomb interactions are completely screened in the Higgs phase, i.e. $Q = 0$, the AB interactions are not. This effect is not a common feature of any screening mechanism, but is specific for screening by a Higgs condensate. For example, Debye screening by a real plasma, or the (sometimes partial) screening related to vacuum polarisation as described by the renormalization group (Gell-Mann-Low β -function), are not of this type. They screen both Coulomb, and AB interactions. Even more peculiar is the screening, related to the Chern-Simons (CS) term [3]. It leads to complete screening of the Coulomb -, and partial screening (by a factor 1/2) of the AB interaction (see for instance Ref. [4], and the text below).

The model in which we discuss these phenomena is the $2+1$ dimensional theory, governed by the Lagrangian

$$\begin{aligned} \mathcal{L} &= \mathcal{L}_{ed}(\mu) + \mathcal{L}_{Higgs} + \mathcal{L}_{matter} \\ &= -\frac{1}{4} F^{\kappa\nu} F_{\kappa\nu} + \frac{\mu}{4} \varepsilon^{\kappa\nu\lambda} F_{\kappa\nu} A_\lambda + |\mathcal{D}_\kappa \Phi|^2 - V(|\Phi|^2) + \mathcal{L}_{matter}. \end{aligned} \quad (1)$$

The first two terms ($\mathcal{L}_{ed}(\mu)$) describe what is known as CS electrodynamics [3]. In the next two terms ($\mathcal{L}_{Higgs}$) a complex scalar field Φ with global $U(1)$ charge Ne is minimally coupled to this gauge theory. In our conventions $\mathcal{D}_\kappa \Phi \equiv (\partial_\kappa + iNeA_\kappa)\Phi$. To proceed, we assume that the potential $V(|\Phi|^2)$ takes the form

$$V(|\Phi|^2) = \frac{\lambda}{4} (|\Phi|^2 - v^2)^2, \quad (2)$$

which spontaneously breaks the gauge symmetry $U(1) \rightarrow Z_N$, and leads to a condensate of the charged Higgs field Φ . In this so-called Higgs phase the length scale is set by $1/M_H$, with $M_H = \sqrt{2\lambda} \langle |\Phi| \rangle = \sqrt{2\lambda} v$ the mass

of the Higgs boson. We have introduced additional charged matter fields ($\mathcal{L}_{matter}$), in order to be able to discuss all conceivable charge sectors in the Higgs phase. We do not further specify these matter fields. We only assume that they are very massive, so that we can discuss the associated global $U(1)$ charges, denoted by q , as external.¹

Let us first recall that pure CS electrodynamics $\mathcal{L}_{ed}(\mu)$ with $\mu \neq 0$ is known to possess only massive particles in the spectrum: one-component massive photons [3] with mass μ . On the other hand, the photons of pure Higgs electrodynamics $\mathcal{L}_{ed}(0) + \mathcal{L}_{Higgs}$ in the Higgs phase are also massive. This time they are endowed with two components, and they carry the mass $M_A = Ne\sqrt{2} |\langle \Phi \rangle| = Ne\sqrt{2}v$. These theories differ radically from massive electrodynamics with explicitly broken gauge invariance

$$\mathcal{L}_{med} = -\frac{1}{4}F^{\kappa\nu}F_{\kappa\nu} + \frac{m^2}{2}A^\nu A_\nu. \quad (3)$$

Despite having only massive particles, the former models allow for the long-range AB interaction [1], which is absent for $\mathcal{L}_{med}$. One may think of the AB interaction in pure CS electrodynamics $\mathcal{L}_{ed}(\mu)$ with $\mu \neq 0$, as being mediated by certain ‘discrete’ degrees of freedom [5], which are close analogues of the by now well known discrete states of the $c = 1$ string [6].

In the Higgs phase of the model $\mathcal{L}_{ed}(0) + \mathcal{L}_{Higgs}$ the AB interaction survives, because the field, which is acquiring mass due to the spontaneous breakdown, is actually the gauge invariant combination

$$\tilde{A}_\kappa \equiv A_\kappa + \frac{1}{Ne}\partial_\kappa \text{Im} \log \langle \Phi \rangle, \quad (4)$$

rather than A_μ itself. Instead of (3), we find in the Higgs phase

$$-\frac{1}{4}F^{\kappa\nu}F_{\kappa\nu} + |\mathcal{D}_\kappa \Phi|^2 \longrightarrow -\frac{1}{4}F^{\kappa\nu}F_{\kappa\nu} + \frac{M_A^2}{2}\tilde{A}^\kappa \tilde{A}_\kappa. \quad (5)$$

Thus $\tilde{A}$ indeed has a finite ($\sim 1/M_A$) correlation length. This does not immediately imply that A should also fall off exponentially. It can instead

¹We take the charges q to be integer multiples of e , i.e. the model (1) corresponds to compact electrodynamics, for instance obtained from some non-abelian model after spontaneous symmetry breaking. In the latter case the topological mass μ would be quantized [3]. In the text we assume however, that μ can take any real value.

remain pure gauge. Such AB fields, which are locally pure gauge, can be globally nontrivial however. This is the case around topological defects of the Higgs condensate of the characteristic size $\sim 1/M_A$, corresponding to magnetic vortices [7] labelled by $\pi_1(U(1)) \simeq Z$. These vortices carry a magnetic flux ϕ which is quantized by the requirement that the Higgs condensate is single-valued outside the core of the vortex, thus ²

$$\phi = \frac{2\pi}{Ne} \times \text{integer}. \quad (6)$$

It is well known that the purely quantummechanical AB interaction [1, 2] leads to non trivial elastic scattering of charges q' , and vortices ϕ , i.e. it gives rise to a diffraction like effect, which is of course observable. ³ The crucial ingredient in the corresponding cross-sections is the phase $\exp iq'\phi$. If there were no magnetic vortices (with their AB fields) in the Higgs phase, electric charges q would be unobservable at *large* distances (of course they can always be seen in scattering processes with high energies $\geq M_A$, i.e. at short distances). But even at *low* energies any two states with $U(1)$ charges q_1 , and q_2 , such that $q_1 - q_2 = Ne \times \text{integer}$ are in fact indistinguishable, because the fluxes which can be used to distinguish them are constrained by Eq. (6). This observation gives rise to the low-energy classification of the superselection sectors of this model by means of (q, ϕ) , with

$$q = e \times (\text{integer mod } N), \quad (7)$$

² Vortices are in fact nothing but 2-dimensional monopoles, however we do not use the word monopole in order to avoid confusion with the monopoles in 3 spatial dimensions. The latter appear in fact as *instantons* in the 2+1 dimensional theory under consideration. These instantons cause transitions with a change of the flux $\Delta\phi = \frac{2\pi}{e}$, and in particular render vortices with $\phi \geq \frac{2\pi}{e}$ unstable [11]. Thus instead of Eq. (6), it is justified to consider a smaller set of fluxes: $\phi = \frac{2\pi}{Ne} \times \text{integer mod } N$ as suggested by the topological characterisation of the fluxes. Even if there were no instantons, two fluxes ϕ_1 , and ϕ_2 with $\phi_1 - \phi_2 = \frac{2\pi}{e} \times \text{integer}$ would be indistinguishable by AB interactions with the charges q , which are integer multiples of e . However, the energy of the fields inside the core of the vortices ϕ_1 , and ϕ_2 is different, and hence are their masses. This may lead to another type of dynamical instability, namely the decay of a vortex with n units of flux into n vortices with a single unit, as it appears in Type II superconductors.

³If the explicit gauge symmetry breaking term $\frac{m^2}{2}A^2$ in (3) is introduced with $1/m \gg 1/M_A$, then the AB field, which was pure gauge, and carried no energy for vanishing m , will acquire a non-vanishing energy. This implies that vortices will be attracted to each other, and therefore will be confined at distances $\sim 1/m$, forming ‘meson’-, and ‘baryon’-like structures of a vortex anti-vortex pair or N vortices respectively.

$$\phi = \frac{2\pi}{Ne} \times (\text{integer mod } N). \quad (8)$$

Together with its less-trivial generalization to the patterns of symmetry-breaking $G \rightarrow H$ with non-abelian discrete groups H [8], this classification has been investigated in Refs. [2], and [9]-[11].

What remains to be discussed is *why* all the states (q, ϕ) , as listed in Eqs. (7), and (8) can indeed be distinguished by the AB interaction. Let us recall that the generic AB field is a pure gauge $A_\mu = g^{-1}\partial_\mu g$ with $g \in U(1)$. In a non-simply-connected domain, one may have that $\oint A_\mu dx^\mu \neq 0$ along a non-contractable loop, and in fact $\oint A_\mu dx^\mu = \phi$ if the contour goes around a vortex once. The interaction of a point-like charge q' (moving along the worldline γ) with the electromagnetic field can be written as $q' \int_\gamma A_\mu dx^\mu$. In the first-quantization formalism the phase factor arising when the charge is carried around the vortex, therefore equals $\epsilon\{\phi, q'\} = \exp i q' \phi$. However, in the Higgs phase the charge q' is completely screened at distances $\gg 1/M_A$. This means that it is surrounded by a ‘cloud’ with a total electric charge exactly equal to $-q'$. This cloud serves to cancel the contribution of q' to the Coulomb field. From a physical point of view, this leads to a potential embarrassment. The problem being that the cloud will be also carried around the vortex, and consequently it seems, that the factor $\epsilon\{\phi, q'\}$ will be *unobservable*, because it should be multiplied by $\epsilon_{\text{induced}}\{\phi, -q'\} = \epsilon\{\phi, q'\}^{-1}$. By this line of reasoning, we are led to conclude that for a physical (i.e. dressed) charge, the AB effect would be absent in the Higgs phase.

Remarkably enough, the reasoning just given, turns out to be incorrect. While the Coulomb interaction is exponentially damped by the Higgs mechanism, the AB interaction is not. In fact $\epsilon_{\text{induced}} \equiv 1$, and the reason for this is that the (induced) screening charge density is *not* an ordinary electric charge density, but rather the field $-M_A^2 \tilde{A}_0$. This is clear from Gauss’s law, implied by Eqs. (1), (4), and (5)

$$\nabla \cdot \mathbf{E}(\mathbf{x}) = q\delta(\mathbf{x}) - M_A^2 \tilde{A}_0(\mathbf{x}) = q\delta(\mathbf{x}) + q_{\text{induced}}(\mathbf{x}), \quad (9)$$

with $E^i \equiv F^{i0}$. The associated induced current is ⁴

$$j_{\text{induced}}^\kappa = -M_A^2 \tilde{A}^\kappa. \quad (10)$$

⁴It is clear that whenever the modulus of the value of the Higgs condensate is varying, as for example inside the core of a vortex, $M_A = (Ne)^2 \sqrt{2} \chi^2$ is also varying, and can in fact become a function of the coordinates, see Eq. (14) below.

In order for this current to interact with the AB field (produced by some remote vortex), there should be a term in the Lagrangian of the form $-j_{\text{induced}}^\kappa A_\kappa = M_A^2 \tilde{A}^\kappa A_\kappa$. Instead we only encounter the term $\frac{1}{2} M_A^2 \tilde{A}^\kappa \tilde{A}_\kappa$ in Eq. (5). In other words, q_{induced} couples to $\tilde{A}$ rather than to A , and thus does *not* feel AB fields related to remote vortices, which have non-vanishing A -component but strictly vanishing $\tilde{A}$ at large distances. Thus our conclusion is that for a physical particle in the Higgs phase the AB interaction is sensitive to the *unscreened* charge only, while ordinary interactions, like the Coulomb one, are completely screened.⁵ The phase factor associated with the AB interaction of two localized states (q, ϕ) , and (q', ϕ') is therefore given by

$$\epsilon\{(q, \phi), (q', \phi')\} \equiv \epsilon\{\phi, q'\} \epsilon\{\phi', q\} = \exp i(q' \phi + q \phi'). \quad (11)$$

Let us now turn to the situation where the CS term is present. The masses of the electromagnetic fields are modified, and become [12]

$$M_\pm = \sqrt{M_A^2 + \frac{1}{2}\mu^2 \pm \mu \sqrt{M_A^2 + \frac{1}{2}\mu^2}}, \quad (12)$$

where $+$, and $-$ stand for two different components of the photon.⁶ The AB field carries no energy, and is still present at large distances. Gauss's law is affected though, it now reads

$$\nabla \cdot \mathbf{E}(\mathbf{x}) = q\delta(\mathbf{x}) + \mu B(\mathbf{x}) - M_A^2 \tilde{A}_0(\mathbf{x}). \quad (13)$$

The new term, $\mu B \equiv \mu F^1_2$, implies that the magnetic field of a vortex generates an additional electric charge density $\mu B(\mathbf{x})$, the total extra charge being $\int \mu B d^2x = \mu \phi$. This extra charge is added to q , and will therefore be completely screened at distances $\gg 1/M_-$ by the screening charge, corresponding to $-M_A^2 \tilde{A}_0$.

⁵ This is a special feature of charge screening in the Higgs mechanism. The Debye screening mechanism, or the screening related to vacuum polarisation as described by the renormalization group, screen both Coulomb -, and AB interactions. This is inferred from the corresponding effective Lagrangians $-\frac{1}{4}F^{\kappa\nu}F_{\kappa\nu} - \frac{m^2}{2}E^i\frac{1}{\Delta}E^i$, and $-\frac{1}{4}F_{\kappa\nu}\frac{e_0^2}{e^2(\Delta)}F^{\kappa\nu}$ respectively.

⁶ If $M_A = 0$ one of the components becomes massless, but then it completely decouples. CS electrodynamics without spontaneous breakdown of gauge symmetry possesses only a *one*-component photon with mass μ .

The most significant effect of the CS term occurs in the phase factor $\epsilon\{(q, \phi), (q', \phi')\}$. In fact, we may think of every physical state (q, ϕ) as being composed of *three* parts: the point-like global $U(1)$ charge q , the vortex ϕ , and the screening charge $q_{\text{induced}} = -q - \mu\phi$ (all concentrated in the domain of radius $\sim 1/M_-$). In the AB field, produced by some remote vortex ϕ' ,

- (a) q is coupled to the total flux ϕ' ,
- (b) q_{induced} does not couple at all,
- (c) and ϕ is coupled to $\frac{\mu}{2}\phi'$.

This can be seen from the form of the Lagrangian density (1), rewritten in terms of $\tilde{A}$

$$-\frac{1}{4}F^{\kappa\nu}F_{\kappa\nu} + \frac{1}{4}\mu\varepsilon^{\kappa\nu\lambda}F_{\kappa\nu}A_\lambda + (Ne)^2\chi^2\tilde{A}^\lambda\tilde{A}_\lambda - V(\chi^2) + (\partial^\kappa\chi)(\partial_\kappa\chi) + \mathcal{L}_{\text{matter}}, \quad (14)$$

where we substituted $\Phi(x) = \chi(x)e^{i\sigma(x)}$ for the Higgs field, with real valued $\sigma(x)$ and gauge invariant $\chi(x) = |\Phi(x)|$, so $\tilde{A}_\kappa \equiv A_\kappa + \frac{1}{Ne}\partial_\kappa\sigma$. The fluctuations of χ around $\langle \chi \rangle = v$ describe neutral Higgs bosons with mass M_H . Now let us assume that we are in the Higgs phase, i.e. χ takes the vacuum expectation value v everywhere. We first note that $\exp(i\sigma/N)$ is not single-valued in the presence of a vortex, it is therefore not a well defined gauge transformation. For this reason we cannot simply replace A by $\tilde{A}$ in the matter term. This implies statement (a). The fact that $\tilde{A}$, rather than A , appears in the interaction term with the Higgs sector, leads to the statement (b) above. Finally, we observe that the coefficient in front of the CS term differs by a factor of 2 in the Lagrangian (14), and in Gauss's law (13). This difference is the origin of the factor $\frac{1}{2}$ in (c) (see for instance Refs. [4, 11]).

From (a)-(c) it follows that the total effect of the AB interaction of the two states (q, ϕ) , and (q', ϕ') when $\mu \neq 0$ is the phase factor [11]

$$\exp i\{(q + \frac{\mu}{2}\phi)\phi' + (q' + \frac{\mu}{2}\phi')\phi\} = \exp i(q\phi' + q'\phi + \mu\phi\phi'). \quad (15)$$

This non-trivial factor arises in the amplitudes, although the total electric charge Q of every localized state (q, ϕ) , if measured by the Coulomb field at distances $\geq 1/M_-$, vanishes

$$Q \equiv q + \mu\phi + q_{\text{induced}} = 0. \quad (16)$$

In Ref. [11], Eq. (15) formed the starting point for the study of the interchange -, and fusion properties of the states (q, ϕ) in the case where $\mu \equiv pe^2/4\pi$ with $p \in Z$. The underlying symmetry was shown to be the Hopf algebra $D(Z_N)$. A non-vanishing value for μ was reflected by a deformation of this algebra by a non-trivial 3-cocycle on the unbroken discrete group Z_N . The generalisation to non-abelian unbroken groups turns out to be straightforward in this formalism

To conclude, we described a mechanism, which makes the fact that electric -, and magnetic fields are massive both in the Higgs phase, and in the presence of a Chern-Simons term in 2+1 dimensions, consistent with the observability of unscreened charges at large distances through long-range Aharonov-Bohm interactions, which survive in both cases. This mechanism allows for smoothly switching on the Chern-Simons term in the Higgs phase, and also provides a solid basis for the discussion of the statistical properties of the states presented in Refs. [10], and [11].

A.B. and M. de W.P. thank Peter van Driel for useful discussions. A.M. is indebted to the Institute for Theoretical Physics of the University of Amsterdam, and the NIKHEF for their kind hospitality and support.

References

- [1] Y. Aharonov, and D. Bohm, Phys. Rev. **115**, 485 (1959).
- [2] L. M. Krauss, and F. Wilczek, Phys. Rev. Lett. **62**, 1221 (1989); M.G. Alford, J. March-Russell, and F. Wilczek, Nucl. Phys. **B337**, 695 (1990); J. Preskill, and L. M. Krauss, Nucl. Phys. **B341**, 695 (1990).
- [3] J. Schonfeld, Nucl. Phys. **B185**, 157 (1981); S. Deser, R. Jackiw, and S. Templeton, Ann. Phys. [N.Y.] **140**, 37 (1982).
- [4] A. S. Goldhaber, R. MacKenzie, and F. Wilczek, Mod. Phys. Lett. **A4**, 21 (1989).
- [5] Y. Kogan, and A. Morozov, Sov. Phys. JETP **61**, 1 (1985).
- [6] I. Klebanov, and A. Polyakov, Mod. Phys. Lett. **A6**, 3273 (1991); E. Witten, Nucl. Phys. **B373**, 187 (1992).

- [7] H. B. Nielsen, and P. Olesen, Nucl. Phys. **B61**, 45 (1973).
- [8] F. A. Bais, Nucl. Phys. **B170** (FSI), 32 (1980).
- [9] M. G. Alford, and J. March-Russell, Int. J. Mod. Phys. **B5**, 2641 (1991); M. Bucher, Nucl. Phys. **B350**, 163 (1991); M. G. Alford, S. Coleman, and J. March-Russell, Nucl. Phys. **B351**, 735 (1991); K. Li, Nucl. Phys. **B361**, 437 (1991); M. Bucher, K. Lee, and J. Preskill, Nucl. Phys. **B386**, 27 (1992); M. G. Alford, and J. March-Russell, Nucl. Phys. **B369**, 276 (1992); M. G. Alford, K. Lee, J. March-Russell, and J. Preskill, preprint CALT-68-1700 (1991).
- [10] F. A. Bais, P. van Driel, and M. de Wild Propitius, Phys. Lett. **B280**, 63 (1992).
- [11] F. A. Bais, P. van Driel, and M. de Wild Propitius, preprint **ITFA 92-8**, to be published in Nucl. Phys. **B**.
- [12] R. D. Pisarski, and S. Rao, Phys. Rev. **D32**, 2081 (1985).